arXiv:1209.5488v1 [hep-ph] 25 Sep 2012

# An Anatomy of Neutrino Oscillations

W-Y. Pauchy Hwang[1]
*Asia Pacific Organization/Center for Cosmology and Particle Astrophysics,
Institute of Astrophysics, Center for Theoretical Sciences,
and Department of Physics, National Taiwan University, Taipei 106, Taiwan*

(September 24, 2012)

## Abstract

To understand neutrino oscillations in the sense of quantum mechanics or quantum field theory, we describe how to use an off-diagonal (cross-generation) neutrino-Higgs (mass) interaction to simulate oscillations in a natural way. This results in an extra orthogonal $SU_f(3)$ family gauge theory, which may help us to resolve a few outstanding puzzles - the question of why there are only three generations, the question of why the masses of neutrinos are so tiny, and the question of why the dark-matter world is so huge (25%) as compared to the visible ordinary-matter world (5%).

PACS Indices: 12.60.-i (Models beyond the standard model); 12.10.-g (Unified field theories and models).

## 1 Introduction

What are neutrinos? Owing to the fact that they have masses, neutrinos may be point-like Dirac particles, as the typical species as one kind of the building blocks of the Standard Model. It might be a species belonging to the dark-matter world, which is known to be five times bigger than the ordinary-matter world (as described by the minimal Standard Model) - in the sense that they interact with dark-matter particles. Now, we know that neutrinos have tiny masses, "tiny" as compared to the other building blocks of the Standard Model. We also know that neutrinos oscillate.

This is not the first example of oscillations, such as the $K^0 - \bar{K}^0$ system, but this is indeed the first example of oscillations for "point-like" (Dirac) particles themselves. In the composite $K^0 - \bar{K}^0$ system, we could draw the Feynman diagrams that would illustrate the underlying mechanism in terms of the minimal Standard Model. But how do we do this for neutrino oscillations? It is a kind of neutrino-flavor changing interaction, at the level of Standard-Model building blocks, very fundamental indeed.

Neutrino oscillations call for a lepton-flavor-violation (LFV) interaction, which brings a neutrino of a given flavor into that of a different flavor. Thus, neutrino masses and neutrino oscillations may be regarded as one of the most important experimental facts over the last thirty years [1].

Using the language of the renormalizable quantum field theory as example, it would correspond to an off-diagonal, cross-generation matrix in the three-neutrino space. It is bilinear in the Dirac fields, so that the coupling to the Higgs field, with some vacuum expectation value, is the only choice, if we insist on the requirement of renormalizability. This Higgs field is the "family" Higgs field, as it operates in the (lepton) family space.

---

[1] Correspondence Author; Email: wyhwang@phys.ntu.edu.tw; a detailed version of arXiv:1207.6443v1 (hep-ph) 27 Jul 2012

## 2 Lepton-flavor-violating interactions

Neutrino oscillations would be the interactions in which neutrino flavors change in the transition. Taking into account neutrino oscillations is equivalent to introduction of lepton-flavor-violating interactions. Using renormalizable quantum field theory, it has to be off-diagonal bilinear in Dirac fields multiplied by the (family) Higgs field with some expectation vacuum values.

Using accelerators, we could produce the "high-energy" neutrino or antineutrino beam, in the energy greater than $2\,GeV$ (i.e. the $\tau$ mass). Considering the $\bar{\nu}_\mu - p$ scattering (the only high-energy neutrino beam available to us, at this point), a fraction of $\bar{\nu}_\mu$ might convert to $\bar{\nu}_\tau$, thus making it possible to produce $\tau^+$. Similarly, for production of $e^+$ via $\nu_\mu$.

To describe the reaction $\bar{\nu}_\mu(k) + p(p) \to \mu(k') + n(p')$, we introduce

$$\begin{aligned} & < n(p') \mid I_\lambda^{(-)}(0) \mid p(p) > \\ = \quad & i\bar{u}(p')\{\gamma_\lambda f_V(q^2) - \frac{\sigma_{\lambda\eta} q_\eta}{2m_p} f_M(q^2)\} u(p), \qquad (1) \\ & < n(p') \mid I_\lambda^{5(-)}(0) \mid p(p) > \\ = \quad & i\bar{u}(p')\{\gamma_\lambda \gamma_5 f_A(q^2) + \frac{i2M q_\lambda \gamma_5}{m_\pi^2} f_P(q^2)\} u(p), \qquad (2) \end{aligned}$$

with $q_\lambda = (p' - p)_\lambda$ and $2M = m_p + m_n$. Here $f_V(q^2)$, $f_M(q^2)$, $f_A(q^2)$, and $f_P(q^2)$ are, respectively, the vector, weak magnetism, axial, pseudoscalar form factors [2]. The transition amplitude is given by

$$\begin{aligned} & T(\bar{\nu}_\mu p \to \mu^+ n) \\ = \quad & \frac{G_F cos\theta_c}{\sqrt{2}} i\bar{v}_\nu(k)\gamma_\lambda(1+\gamma_5)v_\mu(k') < n(p') \mid \{I_\lambda^{(-)}(0) + I_\lambda^{5(-)}(0)\} \mid p(p) > . \qquad (3) \end{aligned}$$

Thus, we obtain [3, 4]

$$\begin{aligned} & \frac{d\sigma}{d\Omega_\mu}(\nu_\mu + p \to \mu^+ + n) \\ = \quad & \frac{G_F^2 cos^2\theta_c E_\mu^2}{2\pi^2}\frac{E_\mu}{E_\nu}\{[f_V^2 + f_M^2 \frac{q^2}{2m_p} + f_A^2 + \frac{m_\mu^2}{m_\pi^2}(f_p^2 \frac{q^2}{m_\pi^2} - 2f_A f_P)]cos^2\theta_\mu \\ & +2[(f_V + f_M)^2 \frac{q^2}{4m_p^2} + f_A^2(1 + \frac{q^2}{4m_p^2}) - 4\frac{E_\mu}{m_p}(1 + \frac{E_\nu}{m_p} sin^2\frac{\theta_\mu}{2}) f_A(f_V + f_M)] sin^2\frac{\theta_\mu}{2}\} \\ \equiv \quad & \frac{G_F^2 cos^2\theta_c E_\mu^2}{2\pi^2}\frac{E_\mu}{E_\nu} \cdot N_0 \\ = \quad & (1.857 \times 10^{-38} cm^2) \cdot (E_\mu,\, E_\nu\ in\ 3\,GeV) \cdot N_0. \qquad (4) \end{aligned}$$

We are dealing with the large side of the weak-interaction cross sections but the neutrino beams cannot be "managed" in terms of the energy and the intensity. But the point which we are trying to get at is to try to see if $\tau^+$ or $e^+$ could be produced in a "high-energy" $\bar{\nu}_\mu$ beam - that might be relatively easy.

In reality, the Higgs field(s) must belong to some group and the coupling is from a neutrino of a certain flavor to the neutrino of different flavor. Thus, it arises naturally the so-called family gauge theory [5]. It is relatively "safe" since the right-handed neutrinos do not appear at all in the minimal Standard Model. We could make a massive $SU_f(3)$ gauge theory completely independent of the minimal Standard Model, including the particle content. Of course, we could treat $(\nu_{\tau R},\, \nu_{\mu R},\, \nu_{eR})$ as a triplet under this $SU_f(3)$ - so to give rise to a completely-independent family gauge theory. Because the anomaly do not hurt in this case, we could drop the right-handed labels from the neutrinos. This completes the derivation of the family gauge theory [5]. The $SU_f(3)$ is by definition the massive gauge theory - all the involved particles, except the neutrinos, are massive dark-matter particles.

We treat the neutrino triplet $(\nu_\tau,\, \nu_\mu,\, \nu_e)$ $(\equiv \Psi)$ as the $SU_f(3)$ triplet. There are six options in this context. To make the $SU_f(3)$ family gauge theory massive, we may choose two complex scalar triplets $(\Phi_+,\, \Phi_-)$ to make all eight gauge bosons massive. To make the mass term, we have the option to write

$$\eta \cdot \frac{1}{2} \cdot (i\bar{\Psi} \times (\Phi_+ + \epsilon\Phi_-) \cdot \Psi + h.c.), \tag{5}$$

as advertised. It is an off-diagonal matrix (in $SU_f(3)$). That is, $\nu_e$ would transform into $\nu_\mu$ or into $\nu_\tau$, $\nu_\mu$ would into $\nu_\tau$ or $\nu_e$, and so on. This is interesting in view of neutrino oscillations, since it could be regarded as the underlying interaction (mechanism) for neutrino oscillations (which we are talking about [1]). An oscillation occurs in a way similar to the decay by way of creating a new species plus the vacuum expectation value (or, changing the vacuum). In quantum mechanics or in quantum field theory, this may be so far the only way how an oscillation can occur.

Now, consider the muon-like neutrino converting to a tau-like neutrino and then producing a tau, i.e. $(\bar{\nu}_\mu \rightarrow \bar{\nu}_\tau) + p \rightarrow \tau^+ + n$. Feynman's rules yield

$$\begin{aligned} iT = \quad & (-)\frac{G_F}{\sqrt{2}} \cdot \eta(u_+ + \epsilon u_-) \cdot \bar{v}(k) \cdot \frac{1}{i}\frac{m_2 - i\gamma \cdot k}{m_2^2 + k^2 - i\epsilon} \cdot \\ & \gamma_\lambda(1+\gamma_5)v(k') \cdot < n(p') \mid [I_\lambda^{(-)}(0) + I_\lambda^{5(-)}(0)] \mid p(p) > \\ = \quad & \{\eta(u_+ + \epsilon u_-) \cdot \frac{i}{m_2 + m_1}\} \cdot \frac{G_F}{\sqrt{2}} \cdot \bar{v}(k)\gamma_\lambda(1+\gamma_5)v(k') \cdot \\ & < n(p') \mid [I_\lambda^{(-)}(0) + I_\lambda^{5(-)}(0)] \mid p(p) > . \end{aligned} \tag{6}$$

Here the small off-diagonal coupling and the small propagator are "canceling" each other. Similarly for the process $(\bar{\nu}_\mu \rightarrow \bar{\nu}_e) + p \rightarrow e^+ + n$.

There is one subtle point here. But a word of caution should be made with the couplings. A pair of complex scalar fields, but eight components get absorbed, have four components left - that means that only four off-diagonal couplings could be different from zero. That is, one of three neutrino oscillations is absent depending on how to arrange $(\nu_\tau,\, \nu_\mu,\, \nu_e)$ in the triplet, or which four nonzero vacuum expectation values are to be chosen. If we choose the four non-zero vacuum expectation values to be first two components, which we choose to the $\nu_\tau$ and $\nu_\mu$ components [5], then the transition from $\nu_\mu$ to $\nu_\tau$ would be absent - in any case, one out of the three would be absent. To make this point clear, we could label the vacuum expectation values $u_\pm$ (in the last equation) as $u_\pm^{(1\to 2)}$, if it is necessary. Note that the irrelevant sign in the transition amplitude, such as in the last equation, could be ignored.

**So, we may try out to do the scattering $\bar{\nu}_\mu + p \to \mu^+ + n$ at the energy of, say, $3\,GeV$. It is interesting to see if $\tau^+$ gets to be produced or if $e^+$ could be produced. This would be a way to open up the door into the "family" world. Of course, we have to worry about the backgrounds; that makes it a high-precision experiment.**

Certain LFV processes [1] such as $\mu \to e + \gamma$ [1] and $\mu + A \to A * + e$ are, in fact, closely related to the most cited picture of neutrino oscillations. See Fig.1 for the illustration. In another note [6, 7], it is realized that the cross-generation or off-diagonal neutrino-Higgs interaction may in fact serve as the detailed mechanism of neutrino oscillations, with some vacuum expectation value of the new Higgs field(s), as contrast with the composed system of the $K^0 - \bar{K}^0$ oscillation.



Figure 1: The leading diagrams for $\mu \to e + \gamma$.

We have the outstanding question why there are three generations in the minimal Standard Model and, yet, another outstanding puzzle that the dark-matter world is about five times the visual ordinary-matter world (the latter described by the minimal Standard Model). Besides the role in the minimal Standard Model, neutrinos may be able to tell us something - an orthogonal family gauge theory in the dark-matter world which our neutrinos are capable of talking to (or interacting with).

Indeed, there is room left for something very interesting. Remember that the right-handed neutrinos never enter in the construction of the minimal Standard Model [4]. The message that the right-handed neutrinos seem to be "unwanted" could be telling us something. Now, the fact that neutrinos have tiny masses suggests that "more naturally" they would be four-component Dirac particles, and unlikely to be the two-component Majorana particles.

We illustrate the point further by calculating the golden lepton-flavor-violating decay $\mu \rightarrow e + \gamma$. We show in Figures 1(a), 1(b), and 1(c) three leading basic Feynman diagrams. Here the conversion of $\nu_\mu$ into $\nu_e$ is marked by a cross sign and it is a term from our off-diagonal interaction given above with the Higgs vacuum expectation values $u_+$ and $u_-$. Here the Higgs masses are assumed to be very large, i.e., greater than a few $TeV$, as in $SU_f(3)$. The only small number (coupling) is $\eta$, explaining the tiny masses of neutrinos.

Using Feynman rules from Wu and Hwang [4], we write, for Fig. 1(a),

$$\begin{aligned}
\frac{1}{(2\pi)^4}\int d^4q \cdot \bar{u}(p',s')\cdot \quad & i\cdot \frac{1}{2\sqrt{2}}\frac{e}{sin\theta_W}\cdot i\gamma_\lambda(1+\gamma_5) \\
\cdot\frac{1}{i}\frac{m_2 - i\gamma\cdot q}{m_2^2+q^2-i\epsilon}\cdot \quad & i\cdot i\eta(-)(u_+ + \epsilon u_-)\cdot\frac{1}{i}\frac{m_1 - i\gamma\cdot q}{m_1^2+q^2-i\epsilon} \\
\cdot i\cdot\frac{1}{2\sqrt{2}}\frac{e}{sin\theta_W}\cdot \quad & i\gamma_{\lambda'}(1+\gamma_5)\cdot u(p,s) \\
\cdot\frac{1}{i}\frac{\delta_{\lambda'\mu}}{M_W^2+(p-q)^2-i\epsilon}\cdot\frac{\epsilon_\sigma(k)}{\sqrt{2k_0}}\cdot \quad & \Delta_{\sigma\mu\nu}\cdot\frac{1}{i}\frac{\delta_{\nu\lambda}}{M_W^2+(p-q-k)^2-i\epsilon},
\end{aligned}$$

with $\Delta_{\sigma\mu\nu} = (-ie)\{\delta_{\mu\nu}(-k-p-q)_\sigma + \delta_{\nu\sigma}(p-q+p-q-k)_\mu + \delta_{\sigma\mu}(-p+q+k+k)_\nu\}$.

On the other hand, Feynman rules yield, for Fig. 1(b),

$$\begin{aligned}
\frac{1}{(2\pi)^4}\int d^4q \cdot \bar{u}(p',s')\cdot \quad & i\cdot \frac{1}{2\sqrt{2}}\frac{e}{sin\theta_W}\cdot i\gamma_\lambda(1+\gamma_5) \\
\cdot\frac{1}{i}\frac{m_2 - i\gamma\cdot q}{m_2^2+q^2-i\epsilon}\cdot \quad & i\cdot i\eta(-)(u_+ + \epsilon u_-)\cdot\frac{1}{i}\frac{m_1 - i\gamma\cdot q}{m_1^2+q^2-i\epsilon} \\
\cdot i\cdot\frac{1}{2\sqrt{2}}\frac{e}{sin\theta_W}\cdot \quad & i\gamma_{\lambda'}(1+\gamma_5)\cdot \\
\cdot\frac{1}{i}\frac{\delta_{\lambda\lambda'}}{M_W^2+(p'-q)^2-i\epsilon}\cdot\frac{1}{i}\frac{m_\mu - 1\gamma\cdot p'}{m_\mu^2+p'^2-i\epsilon}\cdot \quad & i(-i)e\gamma_\sigma\cdot\frac{\epsilon(k)}{\sqrt{2k_0}}.u(p,s),
\end{aligned}$$

and a similar result for Fig. 1(c).

The four-dimensional integrations can be carried out, via the dimensional integration formulae (e.g. Ch. 10, Wu/Hwang [4]), especially if we drop the small masses compared to the W-boson mass $M_W$ in the denominator. In this way, we obtain

$$\begin{aligned}
iT_a = \frac{G_F}{\sqrt{2}} \quad & \cdot\eta(u_+ + \epsilon u_-)\cdot(m_1+m_2)\cdot(+2i)\frac{e}{(4\pi)^2} \\
& \cdot\bar{u}(p',s')\frac{\gamma\cdot\epsilon}{\sqrt{2k_0}}(1+\gamma_5)u(p,s). \qquad (7)
\end{aligned}$$

It is interesting to note that the wave-function renormalization, as shown by Figs. 1(b) and 1(c), yields

$$\begin{aligned}
iT_{b+c} = \frac{G_F}{\sqrt{2}} \quad & \cdot\eta(u_+\epsilon u_-)(m_1+m_2)\cdot(+2i)\frac{e}{(4\pi)^2}\cdot\{\frac{p'^2}{m_\mu^2+p'^2}+\frac{p^2}{m_e^2+p^2}\} \\
& \cdot\bar{u}(p',s')\frac{\gamma\cdot\epsilon}{\sqrt{2k_0}}(1+\gamma_5)u(p,s), \qquad (8)
\end{aligned}$$

noting that $p^2 = -m_\mu^2$ and $p'^2 = -m_e^2$ would make the contribution from Figs. 1(b) and 1(c) to be the same as that from Fig. 1(a).

In a normal treatment, one ignores the wave-function renormalization diagrams 1(b) and 1(c) in the treatment of the decays $\mu \to e + \gamma$, $\mu \to 3e$, and $\mu + A \to e + A^*$.

Comparing this to the dominant mode $\mu \to e\bar{\nu}_e\nu_\mu$ [4], we obtain the branching ratio easily. Even though the decay rate for $\mu \to e + \gamma$, as obtained here, would be of the order $(m_{neutrino}/m_\mu)^4$, which is rather small, it is much larger than order $(m^2/M_W m_\mu)^2$ as often quoted in the literature.

The off-diagonal mass matrix would be modified by the self-energy diagram since the neutrinos form a triplet under $SU_f(3)$. It is presumed that these self-energy diagrams, after the ultraviolet divergences get subtracted, lead to masses of the right order. If the off-diagonal mass matrix is diagonalized alone, the three roots would be two negative and one positive, adding up to zero. This seems like one ordering in the masses of neutrinos - one up and two downs.

In addition to the golden decay $\mu \to e + \gamma$ and the "golden" scattering $(\bar{\nu}_\mu \to \bar{\nu}_\tau) + p \to \tau^+ + n$, one may consider the decay $\pi^0 \to \mu + e$. We would move to a field that would be called "family physics" which deals with baryon-number-conserving lepton-flavor-changing reactions or decays.

Besides the golden decays $\mu \to e + \gamma$ (much too small), $\pi^0 \to \mu + e$, and neutrino oscillations (already observed), we have to decide, to what extent, violation of the $\tau - \mu - e$ universality. As the baryon asymmetry is sometime attributed to the lepton-antilepton asymmetry, the current scenario for neutrino oscillations [1] appears to be inadequate. If we take the hints from neutrinos rather seriously, there are so much to discover, even though the minimal Standard Model for the ordinary-matter world remains to be intact.

## 3 Discussions

Under the assumption that, in the dark-matter world, the dark-matter particles are also species in the extended Standard Model, most of reactions happening among dark-matter particles, even involving neutrinos, cannot be detected in the ordinary-matter world. We could argue that the "minimum" extended Standard Model would be the extended Standard Model to be based on the group $SU_c(3) \times SU_L(2) \times U(1) \times SU_f(3) \times SU_R(2)$ [7], provided that why there are three generations of fermions could be explained and the missing right-handed sector would eventually be coming back [8]. This "minimal extended" Standard Model would be the most natural choice of all the models.

In a slightly different context [9], I proposed to work with two working rules: "Dirac similarity principle", based on eighty years of experience, and "minimum Higgs hypothesis", from the last forty years of experience. By "Dirac similarity principle", all the fermions of the extended Standard Model are point-like Dirac particles and this has important implication for neutrinos. Using these two working rules, the extended model mentioned above becomes rather unique - so, it is so much easier to check it against the experiments. Here we notice that, before the two working rules are adopted, there are so many options (particularly in connection with the Higgs sector) to work out in the left-right models [8].

We would be curious about how the dark-matter world looks like, though it is difficult to verify experimentally. The first question would be if the dark-matter world, 25 % of the current Universe (in comparison, only 5 % in the ordinary matter), would clusterize to form the dark-matter galaxies, maybe even before the ordinary-matter galaxies. The dark-matter galaxies would then play the hosts of (visible) ordinary-matter galaxies, like our own galaxy, the Milky Way. Note that a dark-matter galaxy is by our definition a galaxy that does not "feel" any ordinary strong and electromagnetic interactions (as in our visible ordinary-matter world). This fundamental quest deserves some thoughts, for the structural formation of our Universe.

Of course, we should remind ourselves that, in our ordinary-matter world, those quarks can aggregate in no time, to hadrons, including nuclei, and the electrons serve to neutralize the charges also in no time; then atoms, molecules, complex molecules, and so on. The early stage of "aggregation" does not involve the gravitational force, and so much faster than those by the gravitational force. This early stage of aggregation serves as the "seeds" for the clusters, and then stars, and then galaxies, maybe in a time span of $1\,Gyr$ (i.e., the age of our young Universe). The aggregation caused by strong and electromagnetic forces is fast enough to help giving rise to galaxies in a time span of $1\,Gyr$. On the other hand, the seeded clusterings might proceed with abundance of extra-heavy dark-matter particles such as familons and family Higgs, all greater than a few $TeV$ and with relatively long lifetimes (owing to very limited decay channels). So, further simulations on galactic formation and evolution may yield clues on our problem.

Finally, coming back to the fronts of particle physics, neutrinos, especially the right-handed neutrinos, might couple to the particles in the dark-matter world. Neutrino oscillations and the family "symmetry" are hints for us to enter the new world. Any further investigation along this direction would be of utmost importance. In addition, it may shed light on the nature of the dark-matter world.

This research is supported in part by National Science Council project (NSC 99-2112-M-002-009-MY3). We wish to thank the authors of the following books [4] for thorough reviews of the minimal Standard Model.

# References

[1] Particle Data Group, "Review of Particle Physics", J. Phys. G: Nucl. Part. Phys. **37**, 1 (2010).

[2] H. Primakoff, in "*Muon Physics*", edited by V.W. Hughes and C.S. Wu (Academic, New York, 1975), Vol. II, p.3.

[3] W-Y. Pauchy Hwang, "*Introduction to Particle Physics*" (Lecture notes), unpublished.

[4] Ta-You Wu and W-Y. Pauchy Hwang, "Relatistic Quantum Mechanics and Quantum Fields" (World Scientific 1991); Francis Halzen and Alan D. Martin, "Quarks and Leptons" (John Wiley and Sons, Inc. 1984); E.D. Commins and P.H. Bucksbaum, "Weak Interactions of Leptons and Quarks" (Cambridge University Press 1983). This was "my" early list of the textbooks on the "Standard Model".

[5] W-Y. Pauchy Hwang, Nucl. Phys. **A844**, 40c (2010); W-Y. Pauchy Hwang, International J. Mod. Phys. **A24**, 3366 (2009); the idea first appeared in hep-ph, arXiv: 0808.2091; talk presented at 2008 CosPA Symposium (Pohang, Korea, October 2008), Intern. J. Mod. Phys. Conf. Series **1**, 5 (2011); plenary talk at the 3rd International Meeting on Frontiers of Physics, 12-16 January 2009, Kuala Lumpur, Malaysia, published in American Institute of Physics 978-0-7354-0687-2/09, pp. 25-30 (2009).

[6] W-Y. Pauchy Hwang, arXiv:1207.6443v1 [hep-ph] 27 Jul 2012.

[7] W-Y. Pauchy Hwang, arXiv:1207.6837v1 [hep-ph] 30 Jul 2012; talk presented at the 5th International Symposium on Symmetries in Subatomic Physics, Groningen, Netherland, June 2012; in ”Hyperfine Interactions”, to be published.

[8] J.C. Pati and A. Salam, Phys. Rev. **D10**, 275 (1974); R.N. Mohapatra and J.C. Pati, Phys. Rev. **D11**, 566 (1975); **D11**, 2559 (1975).

[9] W-Y. P. Hwang, arXiv:11070156v1 (hep-ph, 1 Jul 2011), Plenary talk given at the 10th International Conference on Low Energy Antiproton Physics (Vancouver, Canada, April 27 - May 1, 2011), to be published.